# Doc Ref. FP4 Appl. No. 10/593,627

#### (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

### (19) World Intellectual Property Organization International Bureau





### (43) International Publication Date 24 July 2003 (24.07.2003)

### PCT

### (10) International Publication Number WO 03/059388 A1

- (51) International Patent Classification7: A61K 47/02, A61P 9/14
- (21) International Application Number: PCT/IS03/00002
- (22) International Filing Date: 14 January 2003 (14.01.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 6233 15 January 2002 (15.01.2002)
- (71) Applicant (for all designated States except US): DELTA HF. [IS/IS]; Reykjavikurvegi 78. IS-220 Hafnarfjordur (IS).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): EYJOLFSSON, Reynir [IS/IS]; Eyrarholti 6, IS-220 Hafnarfjordur (IS).
- (74) Agent: A & P ARNASON; Efstaleiti 5, IS-103 Reykjavik

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN. YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH. GM. KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM). European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE. ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Declaration under Rule 4.17:

of inventorship (Rule 4.17(iv)) for US only

#### Published:

with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: FORMULATIONS OF QUINAPRIL AND RELATED ACE INHIBITORS

(57) Abstract: Stable formulations of ACE-inhibitors compounds such as quinapril can be produced with the use of excipients comprising a basic compound, preferably an alkali-oralkaline earth metal carbonate, and an insoluble alkaline-earth metal carbonate, and an insoluble alkaline-earth metal hydrogen phosphate. Tablets of such formulations have good storage stability, dissolution characteristics, and the formulations are suitable for use in drug combinations.

WO 03/059388

PCT/IS03/00002

## Formulations of Quinapril and related ACE inhibitors

#### FIELD OF THE INVENTION

5 The present invention is within the field of pharmaceutical formulations of ACE (Angiotensin Converting Enzyme) inhibitors, specifically formulations of quinapril and structurally related compounds.

### 10 TECHNICAL BACKGROUND AND PRIOR ART

Many of the compounds useful as ACE (Angiotensin Converting Enzyme) inhibitors such as as quinapril and structurally related compounds, are prone to degradation. Specifically, such compounds can degrade via (i) cyclization via internal nucleophilic reaction to form substituted diketopiperazines, (ii) hydrolysis of the side-chain ester group, and (iii) oxidation to form products having often unwanted coloration.

Certain stabilizing compositions and formulations of such compounds have been suggested and utilized in the prior art.

20

15

EP 317878 suggests coating an active compound of this type with a polymeric protective coating, or mixing the compound with a physiologically tolerated buffer which ensures that a pH in the weakly acid to weakly alkaline range is set up in a pharmaceutical formulation in the presence of moisture, or both.

25

- The Dictionnaire Vidal (1985) Cahier Complémentaire (p. 10, left col.) describes an enalapril maleate drug named RENITEC which as excipients contains lactose and sodium hydrogen carbonate.
- 30 EP 280999B1 suggests similar compositions using alkali or alkaline earth metal carbonates and saccharides as stabilizers for these compounds. Comparative examples therein (see, e.g. Example D) test a formulation with 5.4 mg quinapril hydrochloride, 88.4 mg magnesium carbonate, 5.2 mg gelatin and 1.0 mg magnesium stearate. The formulation, however, shows substantial degradation as compared with lactose-stabilized formulations and is therefore not useful as a practical pharmaceutical formulation.

Said ACE inhibitor compounds have varied sensitivity and specific formulations need to be tested and optimized for different compounds. Consequently, alternative solutions providing stable formulations will be appreciated.

5

10

It has now been surprisingly discovered that useful, stable formulations can be produced with use of excipients comprising a basic compound, preferably an alkali or alkaline-earth metal carbonate, and an insoluble alkaline-earth metal hydrogen phosphate is further used as a preferred filler substance. Surprisingly, a saccharide compound for stabilization is not needed in such formulations.

The pH of the formulations of the present invention is dominated by the basic stabilizer.

Tablets of such formulations have good storage stability, dissolution characteristics, and the formulations are suitable for use in drug combinations.

In one embodiment, 45% of magnesium carbonate is mixed with 3.9% quinapril hydrochloride with the addition of 33% calcium hydrogen phosphate (CaHPO<sub>4</sub>). Additional specific embodiments and experimental stability test results are disclosed in the Detailed description below and enclosed Examples.

### **DETAILED DESCRIPTION**

25

20

The pharmaceutical formulation of the present invention comprises 0.5 - 50 wt%, such as about 1-25 wt%, including about 1-15 wt% of a compound of formula I:

$$R^{1}$$
 $R^{2}$ 
 $R^{3}$ 
 $R^{4}$ 
 $R^{4}$ 
 $R^{4}$ 

wherein  $R^1$  is hydrogen or alkyl having one to five carbon atoms;  $R^2$  is hydrogen or  $C_1$ - $C_4$  alkyl or the group

(H<sub>2</sub>C) n N B in which A and B independently denote hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl and n is 1-4; R³ and R⁴ together with the atoms they are connected to form a heterocyclic, mono-, di-, or tricyclic ring system which is optionally substituted with C<sub>1</sub>-C<sub>4</sub> alkoxy; R⁵ is methyl or phenyl; or any pharmaceutically acceptable salt thereof; 5 – 90 wt% of an alkali or alkaline earth metal carbonate, such as in the range of about 10 – 90 wt%, including the range of about 15 – 75 wt% and preferably in the range of about 25 – 75 wt%, such as about 25 – 50 wt% or about 30 – 50 wt%; 5 – 90 wt% including the range of about 10 – 90 wt% of an alkali or alkaline earth metal salt of hydrogen phosphate which is preferably an insoluble alkaline earth

5

10

15

20

25

30

35

metal salt of hydrogen phosphate which is preferably an insoluble alkaline-earth metal salt of hydrogen phosphate, such as in the range of about 15 – 75 wt%, such as the range of about 20 – 60 wt%, and preferably in the range of about 25 – 50 wt%, or in the range of 15 – 30 wt%;

with the provisio that the formulation does not contain a substantial amount of a saccharide compound.

In this context a substantial amount of a saccharide compound is meant to include any amount that would generally be considered to have a stabilizing effect on the active compound, such as more than about 10 wt%, and more preferably including an amount which is more than about 5 wt%, or even more preferably including any amount of a saccharide compound which is more than about 2 wt%.

The alkaline earth metal carbonate may suitably be selected from magnesium carbonate, sodium hydrogen carbonate and sodium carbonate.

In preferred embodiments, the amount of the alkaline earth metal carbonate is at least the equivalent of the amount of the active compound of formula I, such as e.g. at least about twice the equivalent, or at least about three times the equivalent of the amount of the active compound.

The term equivalent in this context refers to the conventional ionic equivalent term, one equivalent of a substance participating in a neutralization reaction is that amount of a substance that either contributes or consumes 1 mol of hydrogen ions in that reaction. I.e. for a monoacidic compound such as ramipril and a monobasic alkaline compound such as NaHCO<sub>3</sub>, the equivalent ratio of the compounds is the same as

WO 03/059388 PCT/IS03/00002

the molar ratio; for a diacidic compound such as quinapril HCl stabilized with NaHCO<sub>3</sub>, one equivalent of NaHCO<sub>3</sub> equals two moles of NaHCO<sub>3</sub>.; and likewise for a diacidic compound and dibasic allkaline compound such as Na<sub>2</sub>CO<sub>3</sub>, the equivalent ratio is again the same as the molar ratio. (See, e.g. Skoog, West, Holler Fundamentals of Analytical Chemistry 5th Ed., Saunders College Publishing, NY, 1988).

As mentioned, the pH of the formulations are dominated by the basic stabilizing excipient, i.e., the alkali or alkaline-earth metal carbonate.

10

5

The ACE inhibitor compound is generally selected from enalapril, delapril, lisinopril, moexipril, perindopril, quinapril, ramipril, trandolapril, and pharmaceutically acceptable salts thereof. In particular, stable formulations of quinapril or a salt thereof are suitably manufactured according to the invention disclosed herein.

15

In yet further useful embodiments, the formulation of of the present invention further comprises in the range of about 0.5 – 50 wt% of a pharmaceutically active compound selected from the group containing diuretics including hydrochlorothiazide; antitussives including dextromethorphan, dextromethorphan hydrobromide, noscapine, carbetapentane citrate, and chlophedianol hydrochloride; antihistamines including chloepheniramine maleate, phenindamine tartrate, pyrilamine maleate, doxylamine succinate, and phenyltoloxamine citrate; decongestants including phenylephedrine hydrochloride, phenylpropanolamine hydrochloride, pseudoephedrine hydrochloride, ephedrine; and alkaloids such as codeine phosphate, codeine sulfate, and morphine. The suitable amount of a further pharmaceutically active compound such as the above listed depends on the particular compound, i.e. the activity of the compound and its suitable dose, and the dose weight of the pharmaceutical formulation.

### **EXAMPLES**

### Example 1

5 The following materials were combined by the wet granulation method for the manufacture of 5 mg quinapril tablets.

Quinapril hydrochloride	5.4	mg
Magnesium carbonate	63	mg
Calcium hydrogen phosphate anhydrou	ıs 46.4	mg
Starch pregelatinized	21	mg
Croscarmellose sodium	2.8	mg
Magnesium stearate	1.4	mq

15

# Example 2

The following materials were processed by wet granulation for 10/12.5 mg tablets 20 with Quinapril and Hydrochlorothiazide.

Quinapril hydrochloride	10.8	mg
Hydrochlorothiazide	12.5	mg
Magnesium carbonate	49	mg
Calcium hydrogen phosphate anhydrous	42.5	mg
Starch pregelatinized	21	mg
Croscarmellose sodium	2.8	mg
Magnesium stearate	1.4	mg
<b>U</b>		J

# Example 3

35 Stability of the tablets prepared in the Examples 1-2 were tested at 40°C for 9 days.

	Degradation products (%)			
	Quinapril DKP	Quinaprilat		
Example 1	0.2	1.2		
Example 2	0.3	0.6		

### **CLAIMS**

5

10

15

20

1. A pharmaceutical formulation comprising:

a. 0.5 - 50 wt% of a compound of formula I;

wherein  $R^1$  is hydrogen or alkyl having one to five carbon atoms;  $R^2$  is hydrogen or  $C_1$ - $C_4$  alkyl or the group

 $(H_2C)_{\overline{n}}^A$  in which A and B independently denote hydrogen or  $C_1$ - $C_4$  alkyl and n is 1-4;  $R^3$  and  $R^4$  together with the atoms they are connected to form a heterocyclic, mono-, di-, or tricyclic ring system which is optionally substituted with  $C_1$ - $C_4$  alkoxy;  $R^5$  is methyl or phenyl; and pharmaceutically acceptable salts thereof;

b. 5 - 90 wt% of an alkali or alkaline earth metal carbonate;

c. 5 – 90 wt% of an insoluble alkaline-earth metal salt of hydrogen phosphate;

with the provisio that the formulation does not contain a substantial amount of a saccharide compound.

The formulation of claim 1, wherein the alkali or alkaline-earth metal carbonate is selected from magnesium carbonate, sodium hydrogen carbonate and sodium carbonate.

3. The formulation of claim 1, wherein the amount of the alkaline earth metal carbonate is at least the equivalent of the amount of the active compound of formula I.

WO 03/059388 PCT/IS03/00002

- 4. The formulation of claim 1, wherein the compound of formula I is selected from the group containing enalapril, delapril, lisinopril, moexipril, perindopril, quinapril, ramipril, trandolapril, and pharmaceutically acceptable salts thereof.
- 5. The formulation of claim 1 wherein the formulation comprises in the range of about 1 15 wt% of the compound of formula 1.
  - 6. The formulation of claim 1 wherein the formulation comprises in the range of about 25 75 wt% of the alkali or alkaline earth metal carbonate.
  - 7. The formulation of claim 6 wherein the formulation comprises in the range of about 30 50 wt% of the alkali or alkaline earth metal carbonate.
- 8. The formulation of claim 1 wherein the formulation comprises in the range of about 15 75 wt% of the salt of an insoluble alkaline earth metal hydrogen phosphate.
  - 9. The formulation of claim 1 wherein the formulation comprises in the range of about 25 50 wt% of the salt of an insoluble alkaline earth metal hydrogen phosphate.
  - 10. The formulation of any of the preceding claims wherein the compound of formula I is quinapril or a pharmaceutically acceptable salt thereof.
- 11. The formulation of claim 1 further comprising 0.5 50 wt% of a pharmaceutically active compound selected from the group containing diuretics including hydrochlorothiazide; antitussives including dextromethorphan, dextromethorphan hydrobromide, noscapine, carbetapentane citrate, and chlophedianol hydrochloride; antihistamines including chloepheniramine maleate, phenindamine tartrate, pyrilamine maleate, doxylamine succinate, and phenyltoloxamine citrate; decongestants including phenylephedrine hydrochloride, phenylpropanolamine hydrochloride, pseudoephedrine hydrochloride, ephedrine; and alkaloids such as codeine phosphate, codeine sulfate, and morphine.

10

20

### INTERNATIONAL SEARCH REPORT

Interna lication No PCT/IS 03/00002

A CLASSIF IPC 7	FICATION OF SUBJECT MATTER A61K47/02 //A61P9/14		
According to	International Patent Classification (IPC) or to both national classifica	ition and IPC	
B. FIELDS	SEARCHED cumentation searched (classification system followed by classification	numbala)	
IPC 7	Cumentation searched (classification system followed by classification A61K	in symbols)	
	tion searched other than minimum documentation to the extent that su		
	ata base consulted during the international search (name of data bas		
EPO-In	ternal, CHEM ABS Data, WPI Data, EME	BASE, BIOSIS, MEDLINE	<u> </u>
	ENTS CONSIDERED TO BE RELEVANT		1
Category °	Citation of document, with indication, where appropriate, of the rele	evant passages	Relevant to claim No.
Х	WO 99 62560 A (DANIEL JANE ELLEN LAMBERT CO (US); WEISS JAY (US); 9 December 1999 (1999-12-09) page 3, line 20-25 page 7, line 11-26 page 8, line 6 -page 9, line 17; examples 4,5	HARRIS)	1-11
A	EP 0 280 999 A (WARNER LAMBERT CO 7 September 1988 (1988-09-07) the whole document	))	1-11
Furth	ner documents are listed in the continuation of box C.	X Patent family members are lis	ted in annex.
"A" docume conside filing de "L" docume which i citation "O" docume other n "P" docume	ant defining the general state of the art which is not ered to be of particular relevance occument but published on or after the international atte atte of the state of the s	"T" later document published after the or priority date and not in conflict cited to understand the principle of invention. "X" document of particular relevance; it cannot be considered novel or cal involve an inventive step when the "Y" document of particular relevance; it cannot be considered to involve a document is combined with one of ments, such combination being of in the art.  "A" document member of the same pat	with the application but or theory underlying the the claimed invention nnot be considered to e document is taken alone the claimed invention on inventive step when the or more other such docu- byious to a person skilled
Date of the a	actual completion of the international search	Date of mailing of the international	search report
	8 March 2003	1 0. 04. 2003	
Name and m	nailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer  INGRID EKLUND /	' ELY

## INTERNATIONAL SEARCH REPORT

Intormation on patent family members

Internati plication No
PCT/IS 03/00002

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 9962560	Α	09-12-1999	AU	3979399 A	20-12-1999
			BR	9910947 A	06-03-2001
			CA	2330581 A	1 09-12-1999
			CN	1304322 T	18-07-2001
			EΡ	1083931 A	1 21-03-2001
			HU	0102260 A	
			JP	2002516881 T	
			NO	20006148 A	04-12-2000
			NZ	508544 A	25-10-2002
			PL	344586 A	
			TR	200003600 1	2 20-04-2001
			WO	9962560 A	1 09-12-1999
			US	2002161020 A	
			US	6417196 E	9-07-2002
EP 0280999	Α	07-09-1988	US	4743450 A	10-05-1988
	• •	•••	ΑT	84210 7	
			AU		31-05-1990
			ΑU	1130588 A	25-08-1988
			CA	1300510 A	12-05-1992
			DE	3877226	18-02-1993
			DE	3877226 1	
			DK	94088 A	
			EΡ	0280999 A	
			HK	59496 A	
			ΙE	60931 E	
			JP	2619904 E	
			JP	63225322 A	
			NZ	223407 A	
			PH ZA	26279 A 8800382 A	